

Hypertension

This is a consensus guideline for the pharmacological management of hypertension with frailty. This information was developed by the Dalhousie University Academic Detailing Service and the Palliative and Therapeutic Harmonization (PATH) Program.

This guideline is unique in that it focuses equally on when to stop and when to start medications. We recommend stopping antihypertensive medications that are used for the sole purpose of keeping the systolic blood pressure (SBP) below 140 mmHg, although we are unable to make treatment recommendations for frail older adults with previous stroke (see below).

Recommendations

Carefully review the risks and potential, but unproven, benefits of treatment.

Methods for measuring blood pressure

- Decisions about treatment should be based on blood pressure measurements in the seated (not supine) position, while also considering the presence of orthostasis.
- To evaluate orthostasis, measure BP lying, then immediately on standing and after 2 minutes. Ask the patient if they feel lightheaded or dizzy when standing.

Stopping treatment

- If sitting SBP is <140mmHg, medications can be tapered and discontinued to achieve the targets described in the guideline.
- Before discontinuation, consider if the medications are treating additional conditions, such as rate control for atrial fibrillation or symptomatic management of heart failure.

We are unable to make treatment recommendations for frail older adults at high risk for cardiovascular events. In particular, whether or not to discontinue treatment for individuals with a history of previous stroke is uncertain (see rationale: High Risk due to Previous Stroke)

Starting treatment

- Consider starting treatment when SBP is ≥ 160 mmHg.

- Target SBP should be 140 to 160 mmHg while sitting as long as:
 - There is no orthostatic drop to <140 mmHg using the technique described above.
 - There are no adverse effects from treatment that affect quality of life.
 - See above recommendation regarding treatment of high risk individuals with previous stroke.

- In the very frail with short life expectancy, a target SBP of 160 to 190 mmHg may be reasonable.
- The blood pressure target does not need to change when there is a history of diabetes.
- In general, use no more than 2 medications.

Rationale

- Evidence from “drug treatment” trials (ie, trials that randomize patients to different treatment such as comparing placebo to a drug or comparing one drug to another drug) indicates that there is benefit in treating healthy older adults with hypertension. The benefit of treating frail older adults is unknown.
- Major trials enrolled in elderly patients only if their SBP was above 160mmHg. As such, evidence supports initiating pharmacotherapy at a SBP >160 mmHg. None of the randomized controlled “drug treatment trials” involving elderly patients achieved a SBP <140 mmHg in the active treatment group. Therefore, there is no evidence from randomized controlled trials that supports a SBP target of <140 mmHg for the elderly.
- ‘Treat to target’ trials randomize subjects to two different SBP target goals, but the two groups are treated with the same or similar drugs. Two “treat to target” trials of elderly subjects achieved a SBP <140 mmHg, but there were no statistically significant differences in the primary outcome. Thus, “treat to target” studies also fail to support a SBP target of <140 mmHg for the elderly.
- The benefit of adding a third medication to lower blood pressure has not been studied.
- The characteristics of frailty make the potential benefits of strict blood pressure targets even less certain and increase the possibility of harm from adverse drug events. The only study of adults above the age of 80, HYVET, enrolled relatively healthy subjects and excluded individuals with orthostatic hypotension.

Rationale: High risk due to previous stroke

- Most of the studies reviewed above enrolled relatively healthy older adults. Due to limited evidence, it is even more difficult to judge the potential benefit of lowering BP below 140 mmHg when frail individuals have a history of previous stroke compared to the possibility that medications will cause adverse effects (such as weakness, orthostasis, and falls). To consider treatment benefit with frailty, we valued trial outcomes that would impact quality of life. Thus, a relevant outcome would be non-fatal stroke leading to disability. In contrast, the effect of treatment on mortality is difficult to evaluate when there are competing

- causes for death due to frailty, which makes a mortality outcome less meaningful for the frail.
- In the PROGRESS trial⁹, individuals with a mean age of 64 years were treated with perindopril +/- indapamide. The treatment group experienced decreased rates of disabling stroke, with a relative risk reduction of 38% and absolute risk reduction of 1.64% (2.7% vs 4.3%; NNT 61, [95% CI 39-139]) over almost 4 years, compared to placebo. Based on an evaluation of the risk reduction for all strokes (fatal and non-fatal), the relative risk reduction was found to be similar across a range of baseline systolic pressures, but the absolute reduction was greater in the population with a mean blood pressure of 159/94 mmHg compared to the remainder of the population with a mean blood pressure of 136/79 mmHg. This evidence is based on studies of younger patients in relatively good health; the extent to which these results can be extrapolated to older, frail adults is uncertain due to the timeline needed to achieve benefit and the added vulnerability of frailty, which could make treatment with medications riskier.
 - Another study of individuals with previous stroke and mean age of 66.1±8.6 years, PRoFESS,¹⁰ showed no benefit over 2.5 years in the primary outcome of stroke using telmisartan 80 mg daily compared to placebo. This result is concordant with the PROGRESS trial, which failed to demonstrate a statistically significant reduction in stroke risk with single agent treatment.

References

1. Aronow WS, Fleg JL, Pepine CJ, Artinian NT, Bakris G, Brown AS, et al. ACCF/AHA 2011 Expert Consensus Document on Hypertension in the Elderly: A Report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents. *Circulation* 2011 May 31;123(21):2434-506.
2. Musini VM, Tejani AM, Bassett K, Wright JM. Pharmacotherapy for hypertension in the elderly. *Cochrane Database Syst Rev* 2009 Oct 7;(4):CD000028.
3. Mancia G, Laurent S, Agabiti-Rosei E, Ambrosioni E, Burnier M, Caulfield MJ, et al. Reappraisal of European guidelines on hypertension management: a European Society of Hypertension Task Force document. *J Hypertens* 2009 Nov;27(11):2121-58.
4. Ogihara T, Saruta T, Rakugi H, Matsuoka H, Shimamoto K, Shimada K, et al. Target Blood Pressure for Treatment of Isolated Systolic Hypertension in the Elderly. *Hypertension* 2010 Aug 1;56(2):196-202.
5. Principal results of the Japanese trial to assess optimal systolic blood pressure in elderly hypertensive patients (JATOS). *Hypertens Res* 2008 Dec;31(12):2115-27.
6. Mazza A, Ramazzina E, Cuppini S, Armigliato M, Schiavon L, Rossetti C, et al. Antihypertensive Treatment in the Elderly and Very Elderly: Always "the Lower, the Better?". *Int J Hypertens* 2012;2012:590683. Epub;2011 Sep 22.:590683.
7. Angeli F, Reboldi G, Verdecchia P. "The lower the BP the better" paradigm in the elderly: vanished by VALISH? *Hypertension* 2010 Aug;56(2):182-4.
8. Beckett NS, Peters R, Fletcher AE, Staessen JA, Liu L, Dumitrascu D, et al. Treatment of hypertension in patients 80 years of age or older. *N Engl J Med* 2008 May 1;358(18):1887-98.
9. PROGRESS Collaborative Group. Randomised trial of a perindopril-based blood-pressure-lowering regimen among 6,105 individuals with previous stroke or transient ischaemic attack. *Lancet*. 2001 Sep 29;358(9287):1033-41.
10. Yusuf S, Diener H-C, Sacco RL, Cotton D, Ounpuu S, Lawton WA, et al. Telmisartan to prevent recurrent stroke and cardiovascular events. *N. Engl. J. Med.* 2008 Sep 18;359(12):1225-37.